



## WHAT IS RBD?

REM Behaviour Disorder (RBD) happens during sleep. It can lead to talking and shouting as well as thrashing and hitting out. It can make you move vigorously. You may have vivid, striking dreams as well.

# RBD

## REM Behaviour Disorder



### CONTACT US

T +61 2 9114 0000  
F +61 2 9114 0010  
E [info@woolcock.org.au](mailto:info@woolcock.org.au)

431 Glebe Point Road  
Glebe NSW 2037

[www.woolcock.org.au](http://www.woolcock.org.au)

## REM BEHAVIOUR DISORDER (RBD)

During dreaming, people with RBD are seen to suddenly move or call out. They can look like they are acting out their dreams. The episode usually doesn't go on for long and you might be able to recall what your dream was about when you wake up. The disorder can be a danger to both people with RBD and their bed partners due to the person with RBD responding violently to an imagined attack or because of vigorous movement of their arms and legs.

## WHAT CAUSES RBD?

REM sleep is the dream phase of sleep. This is the time when you have your most structured dreams.

Normally in REM sleep, the muscles should relax totally, so you don't act out your dreams. But if you have RBD, your brain sends the wrong signals to your body muscles so they fail to relax during sleep. This means that when you have dreams you can act these out with bodily movement.

There are two forms of RBD.

- The first often occurs later in life (over 50 years of age) and more often in males. The cause is unknown. RBD may be the first sign of Parkinson's disease and some types of dementia. So people with this form of RBD should be checked for the possibility of the development of these neurological conditions.
- The second form of RBD occurs as a side effect of medicines, such as a high dose of antidepressants. RBD can be made worse if you also have another sleep problem, such as obstructive sleep apnea or periodic leg movements.

## HOW IS RBD DIAGNOSED?

The first step is to work out if the RBD is the side effect of a medication or if there is another obvious cause for it. If no readily identifiable cause can be found, you need to see a sleep specialist for further investigation. He/she will check to see if you have any other sleep problems contributing to the RBD, for example obstructive sleep apnea. Most of the time this assessment will involve an overnight sleep study or Polysomnography. This is done in a hospital/clinic setting and includes measuring the muscle tone in your chin, arms and legs plus video recording you while you sleep.

## HOW DO YOU TREAT RBD?

This depends on the cause. If it appears to be due to a medication side effect, the drug responsible should be stopped if possible. If no specific cause is found there are drugs that can help. If you have been diagnosed with obstructive sleep apnea or another sleep disorder then treatment of this may help the RBD.

If there is a suspicion of other neurological conditions, then a referral to a Neurologist will be recommended.



## WHAT ELSE CAN I DO?

- Adjusting the bedroom to prevent injuries to the patient and partner as necessary.

Make sure there are no sharp corners of furniture by the bed, or glassware nearby.

- If the patient tends to fall onto the floors, it may be worth moving the mattress to the floor.
- Sometimes changing the double bed to twin beds may be necessary.
- Seek advice from a sleep specialist for further assistance.

The Woolcock Institute of Medical Research is a not-for-profit organisation.

If you are interested in further information about becoming involved in our research studies or donations, please visit our website [www.woolcock.org.au](http://www.woolcock.org.au).

Your contribution will make a difference.  
Thank you for your support.

